

**CHEMISTRY 1B – Fall, 2015**  
**FINAL EXAM 1 – VERSION A**

Use Scantron Form SC982-E and select the letter corresponding to the correct answer. Make sure to put **your full name, lab section number, and exam version** (under test no.) on the Scantron Form.

Reference pages, including a periodic table and a page of equations and constants, followed by two pages of scratch paper are provided at the end of the test.

**Part I. Multiple Choice Section.** All Questions have only one correct answer. Each Question is worth 5 points.

1. Which of the following concentration based equilibrium equations correctly corresponds to the following chemical equation:  $\text{CaCO}_3(\text{s}) + \text{H}_2\text{CO}_3(\text{aq}) \rightleftharpoons \text{Ca}^{2+}(\text{aq}) + 2\text{HCO}_3^-(\text{aq})$ ?

a)  $K_C = \frac{[\text{Ca}^{2+}][\text{HCO}_3^-]}{[\text{CaCO}_3][\text{H}_2\text{CO}_3]}$

b)  $K_C = \frac{[\text{Ca}^{2+}][\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]}$

c)  $K_C = \frac{[\text{Ca}^{2+}][\text{HCO}_3^-]^2}{[\text{H}_2\text{CO}_3]}$

d)  $K_C = \frac{[\text{H}_2\text{CO}_3]}{[\text{Ca}^{2+}][\text{HCO}_3^-]}$

2. Given that the  $K_{\text{sp}}$  (sp = solubility constant) for AgCl is  $1.8 \times 10^{-10}$  and the  $K_f$  for formation of  $\text{Ag}(\text{NH}_3)_2^+$  from  $\text{Ag}^+$  and  $\text{NH}_3$  is  $1.7 \times 10^7$ ,  $K$  for:  $\text{AgCl}(\text{s}) + 2\text{NH}_3(\text{aq}) \rightleftharpoons \text{Ag}(\text{NH}_3)_2^+(\text{aq}) + \text{Cl}^-(\text{aq})$  is:

- a)  $1.8 \times 10^{-10}$       b)  $3.1 \times 10^{-3}$       c)  $6.1 \times 10^{-3}$       d) 1.0      e)  $1.7 \times 10^7$

3. For the reaction  $3\text{H}_2(\text{g}) + \text{CO}(\text{g}) \rightleftharpoons \text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g})$ , If the initial partial pressure of  $\text{H}_2$ ,  $\text{CO}$ ,  $\text{CH}_4$  and  $\text{H}_2\text{O}$  are 2.00, 1.00, 0.00 and 0.00 atm, respectively, and the equilibrium partial pressure of  $\text{CH}_4$  is 0.62 atm,  $K_P =$

- a) 0.019      b) 0.27      c) 7.2      d) 52      e) 370

4. Which of the following changes will shift the reaction in problem 3 to the products if you are given that  $\Delta H$  for that reaction is negative?

- a) increase the flask volume      b) increase the temperature  
c) add a catalyst      d) remove some water by condensation  
e) all of the above

5. If  $K_C$  for the reaction  $2\text{Na}_2\text{O}_2(\text{s}) + 2\text{CO}_2(\text{g}) \rightleftharpoons 2\text{Na}_2\text{CO}_3(\text{s}) + \text{O}_2(\text{g})$  is  $6.3 \times 10^4$ , what is the equilibrium concentration of  $\text{CO}_2$  (in M) if the equilibrium concentration of  $\text{O}_2$  is 0.0100 M?

- a)  $1.6 \times 10^{-7}$       b)  $4.0 \times 10^{-4}$       c)  $2.0 \times 10^{-4}$       d) 0.020      e)  $1.3 \times 10^4$

6. A solution has a pH of 4.89.  $[\text{H}^+]$  is:

- a)  $7.8 \times 10^{-10}$       b)  $1.3 \times 10^{-5}$       c)  $7.5 \times 10^{-3}$       d) 0.69      e) 1.3

7. An acid is considered a strong acid if it:
- a) completely dissociates in water                      b) can protonate acetic acid  
 c) is known as a Lewis acid                              d) releases multiple protons per compound  
 e) reacts completely with a strong base
8. What is the pH of a 0.0050 M Ba(OH)<sub>2</sub> solution? (Note: K<sub>w</sub> given in reference material)
- a) 0.01                      b) 2.00                      c) 11.70                      d) 12.00                      e) 12.30
9. What is the pH of a 0.20 M HC<sub>2</sub>H<sub>3</sub>O<sub>2</sub> acid (K<sub>a</sub> = 1.8 x 10<sup>-5</sup>)?
- a) 0.70                      b) 2.72                      c) 4.74                      d) 7.00                      e) 9.26
10. In chlorine containing oxyacids, oxygen is more electronegative than chlorine and stabilizes conjugate anions. Which acid is the strongest?
- a) HClO                      b) HClO<sub>2</sub>                      c) HClO<sub>3</sub>                      d) HClO<sub>4</sub>
11. = BONUS #1. Given K<sub>a</sub> values below, determine which salt solutions will be acidic?  
 a) NaHCO<sub>3</sub>                      b) Na<sub>2</sub>C<sub>2</sub>O<sub>4</sub>                      c) NaHSO<sub>3</sub>                      d) Na<sub>2</sub>HPO<sub>4</sub>                      e) Na<sub>2</sub>SO<sub>4</sub>  
 (Note: K<sub>w</sub> given in reference material)

Acid	K <sub>a1</sub>	K <sub>a2</sub>	K <sub>a3</sub>
H <sub>2</sub> CO <sub>3</sub>	4.5 x 10 <sup>-6</sup>	4.7 x 10 <sup>-11</sup>	NA
H <sub>2</sub> C <sub>2</sub> O <sub>4</sub>	6.0 x 10 <sup>-2</sup>	6.1 x 10 <sup>-5</sup>	NA
H <sub>2</sub> SO <sub>3</sub>	1.6 x 10 <sup>-2</sup>	6.4 x 10 <sup>-8</sup>	NA
H <sub>2</sub> SO <sub>4</sub>	large	1.2 x 10 <sup>-2</sup>	NA
H <sub>3</sub> PO <sub>4</sub>	7.5 x 10 <sup>-3</sup>	6.2 x 10 <sup>-8</sup>	4.2 x 10 <sup>-13</sup>

12. What is the pH of a solution that contains 0.080 M NH<sub>3</sub> and 0.12 M NH<sub>4</sub>Cl?  
 K<sub>b</sub>(NH<sub>3</sub>) = 1.76 x 10<sup>-5</sup>
- a) 4.93                      b) 9.07                      c) 9.25                      d) 9.42                      e) 9.91
13. A chemist has 100.0 mL of a 0.100 M NaC<sub>2</sub>H<sub>3</sub>O<sub>2</sub> solution. 100.0 mL of what solution can be added to the NaC<sub>2</sub>H<sub>3</sub>O<sub>2</sub> solution to make a buffer with pH = pK<sub>a</sub> ± 1? (with pK<sub>a</sub> for HC<sub>2</sub>H<sub>3</sub>O<sub>2</sub>)
- a) 0.050 M HCl                      b) 0.15 M HCl                      c) 0.050 M KC<sub>2</sub>H<sub>3</sub>O<sub>2</sub>                      d) 0.050 M NaOH
14. A chemist creates a new indicator which he calls vomit green that has a pK<sub>a</sub> value of 5.44. It is yellow at pH values less than 5.44 and blue at pH values above 5.44 (with a vomit green shade near 5.44). This best use of this indicator would be for:
- a) titrations of strong bases with strong acids  
 b) titrations of strong acids with strong bases  
 c) titrations of weak acids with strong bases  
 d) titrations of weak bases with strong acids  
 e) titrations of greenish colored solutions

Use the following information to answer questions 15 and 16. A buret is filled with 0.100 M NaOH and it is used to titrate 25.0 mL of HClO, a weak acid of unknown concentration. The titration requires 44.0 mL of NaOH to reach the equivalence point. The  $pK_a$  of HClO is 7.53

15. The original concentration of HClO in the 25.0 mL was:

- a) 0.176 M      b) 0.0568 M      c) 0.114 M      d) 0.0880 M      e) 7.53 M

16. The pH after adding 22.0 mL of NaOH will be:

- a) 0.75      b) 6.77      c) 7.53      d) 8.27      e) 13.00

17. Solid calcium fluoride ( $CaF_2$ ) is added to water to make a saturated solution. Which of the following additions will **decrease** the equilibrium concentration of  $Ca^{2+}$ ?

- a) NaF      b)  $HNO_3$       c) EDTA (forms complex with  $Ca^{2+}$ )      d) more  $CaF_2(s)$

18. Calcium fluoride ( $CaF_2$ ) has a  $K_{sp} = 1.46 \times 10^{-10}$ . Its molar solubility in water is:

- a)  $1.21 \times 10^{-5} M$       b)  $3.32 \times 10^{-4} M$       c)  $6.63 \times 10^{-4} M$       d)  $1.00 \times 10^{-3} M$       e)  $4.67 \times 10^{-2} M$

19. Which of the following reactions leads to a decrease in entropy for the system?

- a)  $H_2O(s) \leftrightarrow H_2O(l)$       b)  $2H_2O(g) + O_2(g) \leftrightarrow 2H_2O_2(g)$   
 c)  $Ag_2CrO_4(s) \leftrightarrow 2Ag^+(aq) + CrO_4^{2-}(aq)$       d)  $H_2O(l) \leftrightarrow H_2O(g)$   
 e)  $Cl_2(g) \leftrightarrow 2Cl(g)$

20. Hydrazine ( $N_2H_4$ ) is used to make rocket fuel and other products but it has positive  $\Delta G_f$  over all temperatures. A strategy to make it would be to:

- a) make from  $N_2$  and  $H_2$ , but using higher temperatures  
 b) make from  $N_2$  and  $H_2$ , but use catalysts  
 c) use a more stable reactant than  $H_2$  (such as  $CH_4$ )  
 d) have the reaction produce another very stable co-product (such as  $H_2O$ )  
 e) have the reaction also produce another unstable product (such as  $N_3$ )

21. Under what temperature regimes will the reaction:  $NH_4NO_3(s) \rightleftharpoons NH_4^+(aq) + NO_3^-(aq)$  ( $\Delta H^\circ = +25.7 \text{ kJ/mol}$ ) be spontaneous?

- a) never      b) high temperature      c) low temperature      d) always

22. The reaction  $H_2(g) + I_2(s) \leftrightarrow 2HI(g)$  has a  $\Delta G^\circ = +3.4 \text{ kJ mol}^{-1}$ . If a system starts with  $P_{H_2} = 0.85 \text{ atm}$  and  $P_{HI} = 0.010 \text{ atm}$  and  $T = 298K$ ,  $\Delta G =$

- a)  $-19 \text{ kJ mol}^{-1}$       b)  $-9.5 \text{ kJ mol}^{-1}$       c)  $-7.6 \text{ kJ mol}^{-1}$       d)  $+3.4 \text{ kJ mol}^{-1}$       e)  $+19 \text{ kJ mol}^{-1}$


23. The  $\Delta G_f^\circ$  values are listed below for molecules in the reaction  $4H_2(g) + 2CO(g) \leftrightarrow C_2H_4(g) + 2H_2O(g)$ . (see reference page for constants)

Compound	$H_2(g)$	$CO(g)$	$C_2H_4(g)$	$H_2O(g)$
$\Delta G_f^\circ$ ( $\text{kJ mol}^{-1}$ )	0	-137.2	68.4	-228.6

The equilibrium constant at 298K is:

- a)  $1.3 \times 10^{-25}$       b) 0.131      c) 1.0      d)  $1.1 \times 10^{20}$       e)  $7.5 \times 10^{24}$

24. The following reaction is an UNBALANCED reaction showing reactants and products of a redox reaction:



When balanced (can assume acidic conditions), the coefficients in front of HClO and Cr (with no fractional coefficients anywhere in the equation) are:

- a) 1 and 1      b) 2 and 1      c) 3 and 1      d) 3 and 2      e) 6 and 2  
, respectively.

25. The oxidation state of C in CH<sub>3</sub>OH is:

- a) +3                      b) +2                      c) +1                      d) 0                      e) -2

For questions 26 to 30, consult the following table of standard reduction potentials as needed.

Reaction	E°(V)
$\text{Cl}_2(\text{g}) + 2\text{e}^- \leftrightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \leftrightarrow 2\text{H}_2\text{O}(\text{l})$	+1.23
$\text{Br}_2(\text{l}) + 2\text{e}^- \leftrightarrow 2\text{Br}^-(\text{aq})$	+1.09
$\text{Ag}^+(\text{aq}) + \text{e}^- \leftrightarrow \text{Ag}(\text{s})$	+0.80
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \leftrightarrow \text{Cu}(\text{s})$	+0.34
$2\text{H}^+(\text{aq}) + 2\text{e}^- \leftrightarrow \text{H}_2(\text{g})$	0.00
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \leftrightarrow \text{Ni}(\text{s})$	-0.23
$\text{Cd}^{2+}(\text{aq}) + 2\text{e}^- \leftrightarrow \text{Cd}(\text{s})$	-0.40
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \leftrightarrow \text{Fe}(\text{s})$	-0.45
$2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \leftrightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83

26. Which of the following reactants can oxidize copper metal (Cu(s)) under standard conditions?

- a) Ag<sup>+</sup>(aq)              b) H<sup>+</sup>(aq)              c) Ni<sup>2+</sup>(aq)              d) Fe<sup>2+</sup>(aq)  
e) all of the above

27. A voltaic cell is made **under standard conditions** by Fe(s) and FeCl<sub>2</sub>(aq) in one half cell and AgNO<sub>3</sub>(aq) and Ag(s) in another half cell (with the two half cells connected by a salt bridge). The voltage measured from the silver (+) to the iron electrode (-) will be:

- a) -0.10 V              b) +0.10 V              c) +0.35 V              d) +1.25 V              e) +2.05 V

28. For the reaction in the voltaic cell made in problem 27, if we want to increase the cell voltage, we can do so by:

- a) increasing the mass of Fe                      b) increasing the mass of Ag  
c) increasing the concentration of FeCl<sub>2</sub>              d) decreasing the concentration of FeCl<sub>2</sub>  
e) using a larger cell with twice the volume (at same concentration) of FeCl<sub>2</sub> and AgNO<sub>3</sub>

29. Which of the metals (in oxidation state of zero) in the list above are oxidized by H<sup>+</sup>?

- a) Ni and Cd              b) Ni only              c) Cd only              d) Cu and Ag              e) Ni, Cd, and Fe

30. BONUS #2 In the following cell, the measured voltage is 0.99 V. Determine x (the concentration of  $\text{AgNO}_3(\text{aq})$  in half of the cell.

cell:  $\text{Ni}(\text{s})|\text{NiCl}_2(\text{aq}, 1.0 \text{ M})||\text{AgNO}_3(\text{aq}, x \text{ M})|\text{Ag}(\text{s})$

- a) 0.21 M                      b) 0.045 M                      c) 0.459 M                      d) 2.2 M                      e) 4.8 M

31. What is a difference between voltaic cells and electrolytic cells?

- a) Voltaic cells have both oxidation and reduction occurring, while electrolytic cells only have reduction occurring  
b) In voltaic cells, oxidation occurs at the anode, while in electrolytic cells, oxidation occurs at the cathode  
c) In voltaic cells, the cells provide electrical energy, while in electrolytic cells, electrical energy is consumed to drive the reaction  
d) In voltaic cells, inert electrodes are used, while in electrolytic cells, metals being oxidized are needed for electrodes  
e) Electrolytic cells can be used for batteries, while voltaic cells can't be used as batteries

32. A redox reaction has a negative  $\Delta G^\circ$  value. This tells us

- a) a negative cell potential is expected under all conditions  
b) a positive cell potential is expected under standard conditions  
c) a K value of less than 1 is expected  
d) this cell would only be useful if it were an electrolytic cell

Questions 33 and 34 involve  $\text{Na}[\text{Cr}(\text{H}_2\text{O})_2(\text{ox})_2]$ . Note that the Latin root for Cr is the same as in English and "ox" stands for  $\text{C}_2\text{O}_4^{2-}$  or oxalate, a bidentate ligand.

33. The name for this complex is:

- a) Sodium diaquadioxalatochromate(III)      b) Sodium biquabioxochromium(III)  
c) Diaquadioxalatochromium(III) sodiate      d) Sodium diaquadioxalatochromate(III)  
e) Sodium bis(dihydromonoxide)dioxalatochromate(III)

34. The oxidation state of the Cr atom is:

- a) +5                      b) +3                      c) +2                      d) +1                      e) 0

35. In coordination complexes, electrons in bonds between ligands and metals almost always come from:

- a) metal s shells                      b) metal d shells                      c) ligand lone pair electrons  
d) ligand inner shell electrons                      e) ligand sigma bonds

36. The coordination complex  $[\text{Ni}(\text{en})_3]^{2+}$  absorbs yellow light (at 560 nm). Calculate  $\Delta$  in kJ/mol. (See reference page for constants)


- a) 214 kJ/mol                      b) 0.214 kJ/mol                      c) 21 kJ/mol                      d) 83,113 kJ/mol

37. The complex  $[\text{Co}(\text{NO}_2)_6]^{4-}$  will have how many unpaired d electrons:

- a) 0                      b) 1                      c) 2                      d) 3                      e) 4

38. Which type of hydrocarbons will have at least two carbons with  $sp$  hybridization?  
a) alkanes          b) alkenes          c) alkynes          d) aromatics


Using the structures shown below, answer the following 2 questions:



39. Which structure is 2,2-dimethylpropane  
a) I                      b) II                      c) III                      d) IV                      e) V
40. Which two structures are isomers (have same molecular formula)?  
a) I and II              b) I and IV              c) IV and V              d) II and IV              e) I and V
41. What is the product of  $CH_2=CHCH_2CH_3$  and  $Cl_2$ ?  
a)  $ClCH_2CH_2CH_2CH_3$                                       b)  $CH_3CClHCH_2CH_3$   
c)  $ClCH_2CClHCH_2CH_3$                                       d)  $CH_2=CHCClHCH_3$   
e)  $CH_2=CHCH_2CH_2Cl$

42. Give the name of the compound to the right.

- a) 1-methyl-3-ethylbenzene                      b) methylphenylethene  
c) 1-ethyl-5-methylbenzene                      d) 1-ethyl-3-methylbenzene



### Constants and Equations

Constants:  $0^{\circ}\text{C} = 273\text{K}$

Universal Gas constant =  $8.314 \text{ J mol}^{-1} \text{ K}^{-1}$

$K_w = 1.0 \times 10^{-14}$  (self hydrolysis constant for water)

$h = \text{Planck's constant} = 6.63 \times 10^{-34} \text{ J}\cdot\text{s}$  and  $c = \text{speed of light} = 3.00 \times 10^8 \text{ m/s}$

Avagadro's number =  $6.02 \times 10^{23}$ .

$F = \text{Faraday's constant} = 96,485 \text{ C/mol e}$

$1 \text{ A (amp)} = 1 \text{ C/s}$

### Equations:

Quadratic Equation – not needed for this test

for  $ax^2 + bx + c = 0$  is  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

### Thermodynamic Equations

$\Delta G_{\text{rxn}} = \Delta G_{\text{rxn}}^{\circ} + RT \ln Q$

$\Delta G^{\circ} = \Delta H^{\circ} - T\Delta S^{\circ}$

### Nernst + related Equations:

$E_{\text{Cell}} = E^{\circ}_{\text{Cell}} - \frac{0.0592}{n} \log Q$

$\Delta G = -nFE^{\circ}$

**Strong ligands:**  $\text{CN}^- > \text{NO}_2^- > \text{en} > \text{NH}_3$     **Weak ligands:**  $\text{H}_2\text{O} > \text{OH}^- > \text{F}^- > \text{Cl}^- > \text{Br}^-$