

# CHEMISTRY 31 EXAM 1

Oct. 5, 2016

NAME \_\_\_\_\_

LAB SECTION # \_\_\_\_\_

## Some Useful Equations and Constants:

Propagation of uncertainty:

Addition/subtraction:

$$y = a + b \text{ or } y = a - b$$

$$S_y = \sqrt{S_a^2 + S_b^2}$$

multiplication/division:

$$y = a \cdot b \text{ or } y = a / b$$

$$\frac{S_y}{y} = \sqrt{\left(\frac{S_a}{a}\right)^2 + \left(\frac{S_b}{b}\right)^2}$$

Exponents:

$$y = a^n$$

$$\frac{S_y}{y} = n \frac{S_a}{a}$$

Note: n = constant  
with no uncertainty

Statistics:

Standard deviation:

$$S = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n - 1}}$$

Grubbs Test:  $G_{\text{calculated}} = \frac{|x_{\text{suspect}} - \bar{x}|}{S}$

F-Test:

$$F_{\text{calc}} = S_1/S_2 \text{ (where } S_1 > S_2 \text{)}$$

Thermodynamics:

$$\Delta G = \Delta G^\circ + RT \ln Q \quad \Delta G^\circ = -RT \ln K \text{ (at equilibrium)} \quad R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}, 0^\circ\text{C} = 273.15 \text{ K}$$

The quadratic equation for  $ax^2 + bx + c = 0$  is  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

**A. Multiple Choice/Fill in the Blank Section.** Only one correct answer for multiple choice questions. (4 points for each question)

1. The SI base units for length, mass, time and amount are:

- a) meter, kilogram, second, and mole
- b) centimeter, gram, second, and mole
- c) meter, pound, minute and dozen
- d) meter, gram, minute, and mole.

2. A concentration is given in parts per million by mass. An equivalent way of expressing this which is true for any solvent is:

- a) (g solute/g solvent)\*10<sup>6</sup>
- b) (g solute)/(10<sup>6</sup> g solution)
- c) (µg solute/kg solution)
- d) (µg solute)/(L solution)

3. A student calibrating a buret in a 0 to 40 mL interval finds that the true volume delivered is 0.07±0.02 mL (mean ± uncertainty) greater than measured volume. When performing a titration with a measured volume of about 40 mL, the student should:

- a) ignore the difference – the measured volume is the true volume
- b) ignore the difference – it is insignificant
- c) subtract 0.07 mL from the measured volume
- d) add 0.07 mL to the measured volume

4. In the calculation  $98.32 - 7.1$ , how many significant figures should be used to express the answer?

- a) 2                      b) 3                      c) 4                      d) 5

5. A researcher has invented a cheaper method to measure glucose in blood. She is interested in comparing the precision of her method compared to a standard method. She measures one blood sample 6 times by her method and the standard method to compare the standard deviations from the two methods. Which test is best to use to compare precision?

- a) case 1 t-test              b) case 2 t-test              c) F-test              d) Grubb's test

6. A new method is tested and found to have small standard deviations, under the 1% relative standard deviation desired. A case 1 t-test indicates significant systematic errors of over 5%. We can conclude this method is:

- a) accurate and precise                      b) precise but not accurate  
c) accurate but not precise                      d) neither accurate nor precise

7. A scientist is using an instrument where response is proportional to concentration. The main purpose of analyzing standards for use with a linear least squares calibration in this case is to:

- a) determine m and b in the equation of a line to be used in the analysis of unknowns  
b) determine if response is related to concentration  
c) determine if the analysis is accurate for unknown compounds  
d) determine if the uncertainty is from the response or the concentration of standards

8. Given that for  $\text{Hg}^{2+} + 2\text{CH}_3\text{CO}_2^- \leftrightarrow \text{Hg}(\text{CH}_3\text{CO}_2)_2(\text{aq})$ ,  $K = 2.8 \times 10^8$  and for  $\text{CH}_3\text{CO}_2\text{H}(\text{aq}) \leftrightarrow \text{H}^+ + \text{CH}_3\text{CO}_2^-$ ,  $K = 1.8 \times 10^{-5}$ ,  $K$  for  $\text{Hg}^{2+} + 2\text{CH}_3\text{CO}_2\text{H}(\text{aq}) \leftrightarrow \text{Hg}(\text{CH}_3\text{CO}_2)_2(\text{aq}) + 2\text{H}^+$ , is:

- a)  $9.1 \times 10^{-2}$               b)  $1.0 \times 10^4$               c)  $2.8 \times 10^8$               d)  $8.6 \times 10^{17}$

9. A reaction requires heat (the reaction container cools) and leads to an increase in disorder. This reaction is:

- a) spontaneous at all T              b) spontaneous at low T              c) spontaneous at high T  
d) never spontaneous

10. For the exothermic reaction  $\text{NH}_3(\text{aq}) + \text{H}^+ \leftrightarrow \text{NH}_4^+$ , which of the following will shift the reaction to the product?

- a) adding water (dilution)                      b) increasing the temperature  
c) adding a strong acid (e.g. HCl)                      d) bubble  $\text{N}_2$  gas through to remove  $\text{NH}_3$  as a gas

**B. Problem Section.** Show all needed calculations to receive full credit. The number of points are shown in parentheses. Use the back side of the page if needed.

1.  $\text{AgNO}_3(\text{aq})$  is being added to precipitate out  $\text{Cl}^-$ . Calculate the volume of 0.40 M  $\text{AgNO}_3$  (in mL) that should be added to 0.61 g of NaCl (formula weight = 58.44 g/mol) so that there is a 10% excess. (8 pts)

2. A metal object is chrome plated by electrical deposition. It is desired to determine the thickness of the chrome plating. The metal object weighs  $142.11 \pm 0.02$  g before plating and  $151.98 \pm 0.02$  g after plating. Given that chrome plating volume = (object surface area)(plating thickness), chrome density =  $7.83 \pm 0.04$  g/cm<sup>3</sup>, and the object surface area =  $61.1 \pm 0.5$  cm<sup>2</sup>, determine the thickness and uncertainty in thickness (in cm). Give to the correct number of significant figures. (14 pts)

Bonus – 3 pts) Which uncertainty contributed the most to the overall uncertainty?

3. A standard of MTBE (methyl tertiary butyl ether) in gasoline, was analyzed 4 times to test a method. The measured values were: 0.81 wt %, 0.68 wt %, 0.73 wt %, and 0.79 %.

a) Using the table below, give the 95% confidence value about the mean using the correct number of significant figures. (8 pts)

Degrees of freedom	3	4	5
t-value (at 95% level)	3.18	2.776	2.571

b) If the true MTBE standard concentration was 0.67 wt %, determine if there was a significant bias in the method at the 95% confidence level. Fully explain your answer. (6 pts)

4. It is desired to measure the concentration of CO emitted from cars. Standards ranging from 25 to 500 ppmv are prepared and analyzed by a CO analyzer giving the following equation for the calibration line:

$$\text{response} = m \cdot \text{concentration} + b$$

where  $m = 0.950 \text{ mV/ppmv}$  and  $b = -12.1 \text{ mV}$

A 16 year-old car gives a response of 214 mV.

a) Calculate the concentration of CO emitted from this car.

b) Is this value reliable (based on where the value falls relative to standards)?

(9 pts)

5. The equilibrium constant for the reaction:  $\text{MnCO}_3(\text{s}) \leftrightarrow \text{Mn}^{2+} + \text{CO}_3^{2-}$  is  $5.3 \times 10^{-10}$  at  $25^\circ\text{C}$ . Determine the value of  $\Delta G^\circ$  in kJ/mol for the reaction. (7 pts)

6. Given that the  $K_{\text{sp}}$  for  $\text{MgF}_2$  is  $3.9 \times 10^{-11}$ , determine the solubility of  $\text{MgF}_2$  in water in mol/L. (8 pts)